

Grazing repellency of methyl anthranilate to snow geese is enhanced by a visual cue

J. Russell Mason^{*†} and Larry Clark[†]

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Denver Wildlife Research Center, c/o Monell Chemical Senses Center, 3500 Market Street, Philadelphia, PA 19104-3308, USA

Methyl anthranilate (Rejex-It AG-36) is formulated as a commercial goose repellent. Frequent reapplications of this product are often necessary, and the cost/application is high ($\geq \$300.00/\text{ha}$). The present experiment tested the possibility that the repellency of methyl anthranilate might be enhanced by the addition of visual cues. Twelve 0.4 ha plots were assigned randomly to three treatment groups. Plots in the first group ($n = 4$) were sprayed with 10% Vapor Guard (an agrochemical adhesive). Plots in the second group ($n = 4$) were treated with a mixture of methyl anthranilate (3.4 kg/ha) and Vapor Guard. Plots in the third group ($n = 4$) were sprayed with a mixture of methyl anthranilate, white paint pigment (titanium oxide, TiO_3 , 0.9 kg/l) and Vapor Guard. From 28 November 1994 to 19 December 1995, all plots were visited at 7 day intervals to collect snow goose (*Chen caerulescens*) droppings along transects. Examination of feces weights/transect meter at 7 days post-treatment showed that both methyl anthranilate formulations reduced goose activity. At 14 and 21 days post-treatment, however, dropping weights were significantly lower in plots treated with methyl anthranilate and TiO_3 than in plots treated with formulated methyl anthranilate alone. These results show that visual cues can enhance the durability of methyl anthranilate repellency.

Keywords: *Chen caerulescens*; damage; goose; grazing; methyl anthranilate; mimicry; Rejex-It; repellent

Populations of Canada geese (*Branta canadensis*) and snow geese (*Chen caerulescens*) are increasing in many areas of North America (Williams and Bishop, 1990; Gauthier and Bedard, 1991). Not surprisingly, reports of crop damage and nuisance complaints also are increasing (Hunt, 1984; Knittle and Porter, 1988; Mason, Clark and Bean, 1993; Mason and Clark, 1994a, 1994b). Because geese are a vector for the transmission of agriculturally important parasites and pathogens (Mason *et al.*, 1993), even farmers without substantial goose damage to their crops complain of geese in their fields.

Existing management strategies include hunting and harassment, planting unattractive cover and lure crops (Owen, 1978, 1990; Gauthier and Bedard, 1991), and using auditory and visual repellents (e.g. Timm, 1983; Conover and Chasko, 1985; Knittle and Porter, 1988; Heinrich and Craven, 1990; Taylor and Kirby, 1990; Mason *et al.*, 1993; Mason and Clark, 1994b). In practice, the use of these techniques is constrained by cost, logistics and effectiveness. Such constraints have stimulated efforts to develop chemical repellents that

are effective, economical and ecologically safe. Methyl anthranilate (CAS 134-20-3), an effective bird repellent (Mason, Adams, and Clark, 1989; Cummings, Mason, Otis and Heisterberg, 1991), was registered in 1994 by the U.S. Environmental Protection Agency (EPA REG. No. 58035-9) as a goose repellent. This substance is the active ingredient in Rejex-It AG-36 (PMC Specialties Group, Cincinnati, OH), and the formulated product is effective against snow geese (Mason and Clark, 1995). However, formulated methyl anthranilate is also expensive ($\geq \$300.00/\text{ha}$ including labor and equipment for application, John Floyd, USDA/APHIS/ADC, pers. comm.), and measurable repellency may require reapplication at 3-4 day intervals. One method to extend the effectiveness of methyl anthranilate may be the use of visual cues. Birds readily associate such cues with unpleasant feeding experiences (Mason, 1988), and they continue to avoid them even in the absence of the causes of underlying unpalatability.

In the present experiment, we compared the goose repellency of formulated methyl anthranilate mixed with a white paint pigment (titanium oxide, TiO_3), to that of formulated methyl anthranilate alone. Titanium oxide was selected as the cue because similar paint pigments (e.g. calcium carbonate) have been used as visual cue additives to aerially applied methiocarb formulations (Dolbeer, Woronecki and Bullard, 1992).

^{*}Author to whom correspondence should be addressed at USDA/APHIS/ADC/NWRC, Predation Ecology and Behavior Project, Utah State University, Logan, UT 84322-5295, USA

[†]Present address: USDA/APHIS/ADC/NWRC, Colorado State University, Engineering Research Center, Foothills Campus, NWRC Room BO5, Fort Collins, CO 80523, USA

Materials and methods

Study sites

We selected 12 0.4 ha study plots on four farms (three plots/farm) in Cumberland and Salem Counties, NJ, USA. The minimum distance between farms was 10 km (range: 10–40 km). All four farms had histories of extensive snow goose grazing (Mason and Clark, 1994b, 1995; Mason *et al.*, 1993).

Our criteria for selection were (a) evidence of goose activity, e.g. feces, footprints, feathers, grazing damage; and (b) agronomic similarity, e.g. planting date, crop, barriers to the wind. Winter wheat (*Triticum aestivum*) was planted in six plots, and Kentucky blue grass (*Poa pratensis*) was planted in the remaining six. We marked the corners of each plot with 0.5 m survey stakes.

Procedure

We randomly assigned the three study plots on each farm to three treatment groups (across farms, $n = 4$ /treatment group), and then established a diagonal transect across each plot. The mean length (\pm standard error) of these transects was 97.4 ± 3.1 m. We removed all debris within 0.3 m of the transect midline on 29 September 1994 with a rake.

Beginning in October 1994, we visited all plots every few days to search for signs of goose activity (i.e. goose droppings along transects). All plots showed signs of goose activity by mid November. On 28–29 November 1994, we again raked the transects to remove goose feces and debris within 0.3 m of the midlines. Treatment formulations were then applied with a Solo Model 410 Backpack Mist Blower. First, we sprayed plots in the control group with an aqueous solution containing 10% Vapor Guard (Miller Chemical Corp., Hanover, PA). Next, we applied formulated methyl anthranilate ($3.4 \text{ kg}^{-1} \text{ ai/ha}$) in a 10% aqueous solution of Vapor Guard to plots in group MA. Finally, we treated the remaining four plots with formulated methyl anthranilate (3.4 kg/ai/ha) in an aqueous solution containing 10% Vapor Guard and 0.9 kg of TiO_3 (Buten Paints, Philadelphia, PA). This group is referred to below as MA/ TiO_3 .

Starting at 7 days post-treatment, we visited all plots at 1 week intervals for 3 weeks. During each visit, we collected all feces within 0.3 m of the midline of each transect. Sampling visits occurred between 0630 and 0900 h, generally prior to the arrival of geese. After collection, droppings were dried for 72 h at 37°C . Dry weights (g) were used as measurements of goose activity within plots (Mason and Clark, 1994b).

To provide an index of the presence or absence of methyl anthranilate, 75×25 mm glass slides were dipped in the treatment formulations on the day of application, and then placed at the midpoint of feces collection transects. Beginning on the day of treatment, four slides were retrieved at weekly intervals for 3 weeks.

Analyses

To determine levels of goose activity as a function of treatment, we calculated mean post-treatment feces

mass/transect meter for each plot. Means were evaluated in a two factor analysis of variance (ANOVA). We considered treatment groups as an independent factor in this analysis and collection dates as a repeated factor. Tukey *post-hoc* tests (Winer, 1962: 193) were used to isolate significant differences among means ($P < 0.05$).

The concentrations of methyl anthranilate residue were determined by standard spectrophotometric techniques (Clark and Shah, 1993; Mason and Clark, 1995). Briefly, each slide was immersed in 100 ml of methanol for 5 days. The methanol samples were sonicated for 30 min and then passed through a $5 \mu\text{m}$ filter to remove encapsulation materials. Filtered solutions were assayed for methyl anthranilate by ultraviolet spectroscopy. Ultraviolet absorbance was measured at 300 nm, and assays of pure methanol served as the control.

Results

There were significant differences between treatment groups ($F = 34.5$; 2,6 df; $P < 0.001$) and among dates ($F = 9.2$; 2,6 df; $P < 0.01$). Also, the interaction between these terms was significant ($F = 6.6$; 4,12 df; $P < 0.005$).

Examination of the main effect for groups showed that mean feces dry mass was lower for MA/ TiO_3 plots than for MA or control plots. Examination of the main effect for dates indicated that feces dry mass was lowest at 7 days post-treatment. Examination of the interaction term revealed the following pattern of effects. At 1 week post-treatment, feces dry masses in MA and MA/ TiO_3 plots were significantly less than in control plots. However, only MA/ TiO_3 plots had a lower dry mass at 2 and 3 weeks post-treatment. Mean feces masses in MA and control plots were approximately the same (Figure 1). Weights in MA/ TiO_3 plots at 2 and 3 weeks were not significantly greater than those recorded at 1 week.

Residue analyses showed that the mean concentration of methyl anthranilate per slide on the day of treatment was 436.5 ± 8.9 mg for MA, and 402.8 ± 6.9 mg for MA/ TiO_3 . At the end of the post-treatment period, the concentration per slide had declined for both groups to 87.7 ± 25.6 mg and 13.1 ± 5.6 mg per slide, respectively.

Discussion and management implications

Goose activity levels (i.e. mean feces/transect meter) are indicative of grazing damage (Mason and Clark, 1994b, 1995). Accordingly, both MA and MA/ TiO_3 reduced damage for 7 days post-treatment. Only MA/ TiO_3 continued to prevent damage throughout the post-treatment period, however, suggesting that the visual cue was a useful additive to the formulation. This difference in effectiveness ($\geq 300\%$) is important, because the cost of methyl anthranilate applications is high.

While it is conceivable that TiO_3 alone conferred some repellency, we believe that this is unlikely. Firstly, pilot tests suggested that TiO_3 alone does not deter grazing by geese (Mason, unpubl. data).




Figure 1. Mean feces mass g/transect meter in plots treated with methyl anthranilate (A636), methyl anthranilate plus titanium oxide (A6-6 + TiO₂) or left untreated

Secondly, in other field experiments, simple applications of white paint pigment (i.e. calcium carbonate) were not avoided (Dolbeer *et al.*, 1992).

We suspect that any visual cue that contrasts against background vegetation would serve at least as well as TiO₃. Colors that serve biological warning functions (so-called aposematic cues) may be the most effective (Mason, 1988). Colored plastic flags might also enhance methyl anthranilate repellency. In fact, flagging may be even more effective than color additives, *per se*, because flags are inherently repellent to geese (Mason and Clark, 1994b, 1995). This hypothesis is readily testable. Finally, an intriguing possibility is that methyl anthranilate/visual cue applications might be coupled with applications of visual cues (pigments, flags) alone. Analogous to automimicry effects (Mason, 1988), the methyl anthranilate/visual cue combinations (i.e. models) may confer significant protection to vegetation treated with the visual cue alone.

Acknowledgements

Funding was provided by US Department of Agriculture Cooperative Agreement No. 12-34-41-0040 (CA) between the Monell Chemical Senses Center and the Denver Wildlife Research Center (DWRC). We thank Stella Farms, Inc., Elmer, NJ and Eastern Fresh Farms, Cedarville, NJ, for the use of their fields. Rejex-It AG-36® was donated by PMC Specialities Group (Cincinnati, Ohio, USA). Use of Rejex-It AG-36 does not imply endorsement by the US Department of Agriculture. Special thanks are extended to Dr R. Herz for a critical review of an early manuscript draft.

References

Clark, L. and Shah, P. S. (1993) Bird repellents: possible use in cyanide ponds. *J. Wildlife Mgmt.* 57, 657-664

Conover, M. R. and Chasko, G. C. (1985) Nuisance Canada goose problems in the eastern United States. *Wildlife Soc. Bull.* 13, 228-233

Cummings, J. L., Mason, J. R., Otis, D. L. and Heisterberg, J. L. (1991) Evaluation of dimethyl and methyl anthranilate as a Canada goose repellent on grass. *Wildlife Soc. Bull.* 19, 184-190

Dolbeer, R. A., Woronecki, P. P. and Bullard, R. W. (1992) Visual cue fails to enhance bird repellency of methiocarb in ripening sorghum. In: *Chemical Signals in Vertebrate, VI* (Ed. by D. Muller-Schwarze, R. Silverstein and D. Doty), pp. 323-330. New York, Plenum Press

Gauthier, G. and Bedard, J. (1991) Experimental tests of the palatability of forage plants in greater snow geese. *J. Appl. Ecol.* 28, 491-500

Heinrich, J. W. and Craven, S. R. (1990) Evaluation of three damage abatement techniques for Canada geese. *Wildlife Soc. Bull.* 18, 405-410

Hunt, R. A. (1984) Crop depredations by Canada geese in east-central Wisconsin. *Proc. Eastern Wildlife Damage Control Conf.* 1, 245-254

Knittle, C. E. and Porter, R. D. (1988) *Waterfowl Damage and Control Methods in Ripening Grain: An Overview*. Washington, DC, U.S. Fish and Wildlife Service, 17 pp

Mason, J. R. (1988) Direct and observational avoidance learning by red-winged blackbirds (*Agelaius phoeniceus*): the importance of complex visual stimuli. In *Social Learning: A Biopsychological Approach* (Ed. by T. Zentall and B.G. Galef), pp. 99-115, Lawrence Erlbaum, Hillsdale, N.J.

Mason, J. R., Adams, M. A. and Clark, L. (1989) Anthranilate repellency to starlings: Chemical correlates and sensory perception. *J. Wildlife Mgmt.* 53, 55-64

Mason, J. R. and Clark, L. (1994a) Use of activated charcoal and other particulate substances as feed additives to suppress bird feeding. *Crop Protect.* 13, 219-225

Mason, J. R. and Clark, L. (1994b) Evaluation of plastic and mylar flagging as repellents for snow geese (*Chen caerulescens*). *Crop Protect.* 13, 531-534

Mason, J. R. and Clark, L. (1995) Evaluation of RejexIt AG36 and Anjan-Activaid as snow goose grazing deterrents. *Crop Protect.* (in press)

Mason, J. R., Clark, L. and Bean, N. J. (1993) White plastic flags repel snow geese (*Chen caerulescens*). *Crop Protect.* 12, 497-500

Owen, M. (1978) Food selection in geese. *Verhandlungen der Ornithologischen Gesellschaft in Bayern* 23, 169-176

- Owen, M.** (1990) The damage-conservation interface illustrated by geese. *Ibis* **132**, 238–252
- Taylor, J. P. and Kirby, R. E.** (1990) Experimental dispersal of wintering Snow and Ross' geese. *Wildlife Soc. Bull.* **18**, 312–319
- Timm, R. M.** (1983) *Prevention and Control of Wildlife Damage*. University of Nebraska Press, Lincoln, NE, 632 pp
- Williams, B. K. and Bishop, R.** (1990) Perspectives on goose management in North America: challenges and opportunities for the 1990s. *Trans. North American Wildlife and Natural Resources Conf.* **55**, 283–285
- Winer, B. J.** (1962) *Statistical Principles in Experimental Design*. McGraw-Hill, New York, 907 pp

Received 27 June 1995

Revised 15 September 1995

Accepted 18 September 1995